



Ladder webs in orb-web spiders: ontogenetic and evolutionary patterns in Nephilidae

MATJAŽ KUNTNER^{1,2*}, SIMONA KRALJ-FIŠER¹ and MATJAŽ GREGORIČ¹¹Institute of Biology, Scientific Research Centre, Slovenian Academy of Sciences and Arts, Novi trg 2, PO Box 306, SI-1001 Ljubljana, Slovenia²Department of Entomology, National Museum of Natural History, Smithsonian Institution, NHB-105, PO Box 37012, Washington, D.C. 20013-7012, USA

Received 18 September 2009; accepted for publication 10 November 2009

Spider web research bridges ethology, ecology, functional morphology, material science, development, genetics, and evolution. Recent work proposes the aerial orb web as a one-time key evolutionary innovation that has freed spider-web architecture from substrate constraints. However, the orb has repeatedly been modified or lost within araneoid spiders. Modifications include not only sheet- and cobwebs, but also ladder webs, which secondarily utilize the substrate. A recent nephilid species level phylogeny suggests that the ancestral nephilid web architecture was an arboricolous ladder and that round aerial webs were derived. Because the web biology of the basalmost *Clitaetra* and the derived *Nephila* are well understood, the present study focuses on the webs of the two phylogenetically intervening genera, *Herennia* and *Nephilengys*, to establish ontogenetic and macroevolutionary patterns across the nephilid tree. We compared juvenile and adult webs of 95 *Herennia multipuncta* and 143 *Nephilengys malabarensis* for two measures of ontogenetic allometric web changes: web asymmetry quantified by the ladder index, and hub asymmetry quantified by the hub displacement index. We define a 'ladder web' as a vertically elongated orb exceeding twice the length over width (ladder index ≥ 2) and possessing (sub)parallel rather than round side frames. Webs in both genera allometrically grew from orbs to ladders, more so in *Herennia*. Such allometric web growth enables the spider to maintain its arboricolous web site. Unexpectedly, hub asymmetry only increased significantly in heavy-bodied *Nephilengys* females, and not in *Herennia*, challenging the commonly invoked gravity hypothesis. The findings obtained in the present study support the intrageneric uniformity of nephilid webs, with *Herennia etruscilla* webs being identical to *H. multipuncta*. The nephilid web evolution suggests that the ancestor of *Nephila* reinvented the aerial orb web because the orb arises at a much more inclusive phylogenetic level, and all intervening nephilids retained the secondarily acquired substrate-dependent ladder web. © 2010 The Linnean Society of London, *Biological Journal of the Linnean Society*, 2010, 99, 849–866.

ADDITIONAL KEYWORDS: Araneae – Araneoidea – *Clitaetra* – *Herennia* – hub displacement – ladder index – *Nephilengys* – *Nephila* – architecture – web allometry.

INTRODUCTION

Spider webs may be viewed as the animal's extended phenotype, or a static manifestation of the spider's behaviour (Eberhard, 1982; Schuck Paim, 2000; Kuntner, Coddington & Hormiga, 2008a). A mass of literature exists addressing all aspects of spider-web biology from architecture (Eberhard, 1975, 1977, 1985, 2007; Coddington, 1986c; Agnarsson & Cod-

dington, 2006), web construction and predatory behaviour (Eberhard, 1981, 1986, 1987, 1990, 1992; Rypstra, 1982; Coddington & Sobrevala, 1987; Blackledge, 1998; Herberstein & Heiling, 1998; Li, 2005; Blackledge & Zevenbergen, 2006; Zschokke *et al.*, 2006), web ecology (Toft, 1987; Higgins, 1992, 2006; Higgins & Buskirk, 1992; Wise, 1993; Tso, 1996; 1998a, b; Agnarsson, 2003; Tso, Jiang & Blackledge, 2007; Kuntner *et al.*, 2008b), web ontogenetic allometric changes (Eberhard, 1985; Japyassu & Ades, 1998; Kuntner *et al.*, 2008b; Kuntner & Agnarsson,

*Corresponding author. E-mail: kuntner@gmail.com

2009), web silk properties and the underlying genetics (Denny, 1976; Hayashi & Lewis, 2000; Hayashi, 2001; Hayashi, Blackledge & Lewis, 2004; Blackledge, Summers & Hayashi, 2005; Blackledge, Swindeman & Hayashi, 2005; Agnarsson *et al.*, 2009; Opell & Schwend, 2009), and web evolution (Shear, 1986; Benjamin & Zschokke, 2003, 2004; Craig, 2003; Zschokke, 2003; Blackledge & Gillespie, 2004; Vollrath & Selden, 2007; Japyassu & Caires, 2008; Eberhard, Agnarsson & Levi, 2008a; Eberhard, Barrantes & Madrigal-brenes, 2008b).

Orb webs of orbicularian spiders are highly efficient traps, and their evolution has taken many different paths, resulting in surprising diversity and variation (Eberhard, 1982; Coddington, 1986b, c; Coddington & Levi, 1991). The orb web has evolved only once and its origin defines Orbiculariae with approximately 11 000 species (Blackledge *et al.*, 2009; Coddington, 1986a–c; Coddington, 1989; Garb *et al.*, 2006). However, the architecture is not evolutionarily stable, having been modified into sheet- or cobwebs, or even abandoned (Griswold *et al.*, 1998). Previous studies interpreted the shift from cribellate webs to aerial orb webs made of adhesive silk as a key innovation (Bond & Opell, 1998; Blackledge *et al.*, 2009). Vertical aerial webs represent an escape from the constraints of substrate-based webbing, and gluey silk is much more efficient and economical than the more primitive, cribellar silk (Blackledge *et al.*, 2009). However, araneoid spider clades that continue to build orb webs with adhesive silk have developed an array of architectures that depart from the archetype, which is the approximately round, aerial web, only touching the substrate via anchor threads (Zschokke, 1999). Some groups have secondarily modified the orb to be more substrate dependent and these construct their webs against tree trunks, other vegetation, rock outcrops, and artificial constructions (Kuntner, 2005, 2006, 2007; Harmer & Framenau, 2008). Such substrate-based orb modifications are sometimes termed ladder webs because they are often or always vertically elongated (Kuntner *et al.*, 2008a, b; Harmer, 2009; Harmer & Herberstein, 2009).

Current phylogenetic evidence (Blackledge *et al.*, 2009; Scharff & Coddington, 1997; Kuntner *et al.*, 2008a) suggests that such specialized orb webs evolved independently in the different araneoid groups: the neotropical araneid *Scoloderus* (Eberhard, 1975), the Australian araneid *Telaprocera* (Harmer & Framenau, 2008; Harmer, 2009; Harmer & Herberstein, 2009), the New Zealand araneid *Cryptaranea* (Forster & Forster, 1985), the New Guinean tetragnathid *Tylorida* (Robinson & Robinson, 1972), and the ancestor to the extant nephilid spiders, where *Clitaetra* and *Herennia* show different, yet homologous ladder webs (Robinson & Lubin, 1979; Kuntner, 2005,

2006; Kuntner *et al.*, 2008a). The convergent evolution of ladder webs appears to be the result of different selective pressures. For example, the extreme aerial ladder in *Scoloderus* is an adaptation for moth catching (Eberhard, 1975; Stowe, 1978). By contrast, there is evidence that some ladder-web spiders are able to adapt their web architecture to the substrate at hand and thus alternate between phenotypes (e.g. *Telaprocera*; Harmer & Herberstein, 2009). Thus, the *Telaprocera* ladder web is not an obligate architecture, but rather a plastic response to the available microhabitat. In the present study, we examine web biology of substrate utilizing spiders of the family Nephilidae, which show remarkable intrageneric uniformity but substantial intergeneric diversity in web form, spanning different levels of vertical elongation (Kuntner *et al.*, 2008a; Kuntner & Agnarsson, 2009). The goals of the study are to assess ontogenetic allometric changes in these webs, define ladderness, and establish whether nephilid ladder webs are a response to evolutionary trends (phylogenetic inertia) or to behavioural plasticity (response to available space).

Parsimony ancestral character reconstruction suggests that ladder shaped orb webs on tree trunks found in known species of the *Clitaetra-Herennia* grade were the ancestral life style of the pantropical clade Nephilidae and have apparently reversed to aerial orb webs in the ancestor of *Nephila* (Kuntner *et al.*, 2008a; Kuntner & Agnarsson, 2009): *Nephila* webs are not constrained by the substrate, and do not resemble a ladder (Harvey, Austin & Adams, 2007; Kuntner *et al.*, 2008a). Kuntner *et al.* (2008b) studied *Clitaetra irenae* Kuntner 2006 and quantified its ontogenetic web changes from orb web to ladder web and the simultaneous hub displacement (HD) towards the top frame in older spiders. They concluded that: (1) increasing levels of ladder-web architecture allow the web size to increase disproportionately more vertically than horizontally; this enables the growing spider to remain on the same tree throughout its life, and (2) the logical explanation for an ontogenetic shift from a central hub in small juveniles towards the eccentricity seen in larger, heavier spiders, is gravity, because predation success of heavier spiders is optimized in webs with hubs displaced above the web centre (Masters & Moffat, 1983). Kuntner & Agnarsson (2009) tested these trends in two previously unstudied *Clitaetra* species and concluded that, although allometric shifts in *Clitaetra episinooides* and *Clitaetra perroti* were less pronounced compared to *C. irenae*, *Clitaetra* nevertheless showed within-genus web uniformity. The same seems to hold for other nephilid genera (Kuntner & Agnarsson, 2009).

With the webs of the relatively basal and derived nephilid genera, *Clitaetra* and *Nephila* already



Figure 1. Juvenile and adult web architectures of *Herennia multipuncta* and *Nephilengys malabarensis* from Singapore: A, *Herennia multipuncta* adult female web ($a = 16\text{cm}$, $b = 72\text{cm}$, $c = 21\text{cm}$) with hub cup (HC) and pseudoradial (P). B, *Herennia multipuncta* third-instar web ($a = 6.5\text{cm}$, $b = 16\text{cm}$, $c = 7.3\text{cm}$) with hub (H) not touching bark and parallel side frames (S). C, *Nephilengys malabarensis* third-instar web ($a = 10.3\text{cm}$, $b = 11.5\text{cm}$, $c = 5.6\text{cm}$) with labelled web measurements. D, *Nephilengys malabarensis* adult female web ($a = 43\text{cm}$, $b = 73\text{ cm}$, $c = 24\text{cm}$) with retreat (R) connected to the hub. a , web width; b , web height; c , hub to top distance.

studied, the focus of the present study is on web biology of the remaining two nephilid genera, *Herennia* and *Nephilengys*. Considering their intervening phylogenetic placement on the nephilid tree, one might expect also intermediate web forms between small ladder webs (*Clitaetra*) and quite enormous aerial webs (*Nephila* reaching 1.5 m in diameter; Kuntner *et al.*, 2008a). We investigated web architectures of juvenile and adult spiders of one species of each genus aiming to establish ontogenetic and macroevolutionary trends across the phylogeny. We tested the ontogenetic web allometry shifts as seen in *Clitaetra* (Kuntner *et al.*, 2008b; Kuntner & Agnarsson, 2009), and predicted that: (1) these arboricolous

spiders will likewise show a significant increase in ladder-shape of their webs through ontogeny and (2) their hub eccentricity will be even more pronounced than that found in *Clitaetra* as a result of gravity because *Herennia*, and especially *Nephilengys* adults are considerably larger than *Clitaetra*. We built our predictions upon our knowledge of adult nephilid web architecture (Kuntner *et al.*, 2008a): adult female *Herennia* are known for extreme ladder webs (Fig. 1) (Kuntner, (2005) and adult female *Nephilengys* are known for substrate-webs with displaced hubs (Fig. 2) (Kuntner, (2007)). Additionally, Japyassu & Ades (1998) reported that juvenile *Nephilengys cruentata*, unlike adults, still make round orbs. Our second




Figure 2. Box plots of measured web parameters for most size classes in *Herennia multipuncta* and *Nephilengys malabarensis*. *P*-values are probabilities for the difference between size classes (Kruskal–Wallis test, above) and probabilities for the correlation between size classes and parameters (Spearman's rank order correlation, below). Using Bonferroni correction, *P* is significant at 0.005 or lower.

prediction is also supported by the evolution of size in nephilid spiders: adult female size (and thus weight) gradually increases through the phylogeny from small bodied *Clitaetra* to evolutionarily gigantic *Nephila* (Kuntner & Coddington, 2009).

MATERIAL AND METHODS

We studied web architectures in *Herennia multipuncta* (Doleschall, 1859) and *Nephilengys malabarensis* (Walckenaer, 1842) in the field in Singapore and Indonesia. Web measurements of 95 *H. multipuncta* and 143 *N. malabarensis* randomly encountered individuals were taken along with notes of their habitat. The third species encountered in the field, but not in statistically sufficient quantities (20 individuals), was *Herennia etruscilla* Kuntner 2005. Considering the intrageneric uniformity of nephilid webs (Kuntner *et al.*, 2008a; Kuntner & Agnarsson, 2009), we considered *H. multipuncta* and *N. malabarensis* to be representative of the genera *Herennia* and *Nephilengys* and only analysed *H. etruscilla* data with the intention to test this trend.

We observed web characteristics and measured habitat and web parameters (see Appendix, Table A1) in finished webs of those spiders that could be reliably identified and sized: habitat type (primary versus secondary forest, artificial constructions), tree species if known, bark type (rough, medium, smooth), tree circumference at the hub, canopy closure (closed, partially, and fully open), hub height from ground, web width (Fig. 1C: a), height (Fig. 1C: b) and distance between top frame and the hub (Fig. 1C: c), side frame curvature (round, subparallel, parallel; Fig. 1B, C), web shape (planar, convex, concave), the presence of a hub-cup (Fig. 1A), pseudoradii (Fig. 1A), and a retreat (Fig. 1D; Kuntner, 2005, 2007; Kuntner *et al.*, 2008a, b). As in our previous studies (Kuntner *et al.*, 2008b; Kuntner & Agnarsson, 2009), we assigned the spiders to estimated size class in the field (size classes 2–8 = estimated instar numbers, where '8' indicates adult females), but also tested the validity of such technique by regressing the size class estimations with real specimen measurements for a subset of the specimens.

The two ratios quantifying web allometry are based on previous studies [Kuntner *et al.* (2008b); Kuntner & Agnarsson (2009)]. The *ladder index* (LI), equivalent to web elongation *sensu* Harmer (2009) and similar to web shape *sensu* Zschokke (1993) and web asymmetry *sensu* Blackledge & Gillespie (2002), is the relative web height, defined as the ratio of web height to web width (b/a), where b = web height and a = web width (Fig. 1A, B, C). LI quantifies web elongation (i.e. the orb versus ladder-web architecture), which is known to increase during ontogeny in *Clita-*

etra (Kuntner *et al.*, 2008b). Previous studies have used this index to describe the properties of ladder webs, although the definition of the 'ladder' has been lacking. We define a ladder web to be at least twice as high as it is wide ($LI \geq 2$) and to possess (sub)parallel, rather than round, side frames. Hub displacement (HD), similar to hub asymmetry *sensu* Blackledge & Gillespie (2002), is web height below hub/total web height using the formula $(b - c)/b$, where b = web height and c = distance from top frame to hub (Fig. 1A, B, C). HD values increase with the hub being displaced towards the top web frame, which is a more logical measure of hub eccentricity compared to hub asymmetry indices with decreasing values (Masters & Moffat, 1983; ap Rhisiart & Vollrath, 1994; Kuntner *et al.*, 2008a). Harmer (2009) studied upper/lower asymmetry in the ladder webs of *Telaprocera* spiders, and emphasized the comparison of upper to lower web area rather than the simple relative position of the hub, as detected by HD.

We analysed within species differences in numeric parameters along size classes by the Kruskal–Wallis test and their relationships using Spearman's rank order correlation. The intra- and interspecific differences in categorical parameters were tested by contingency tables and chi-square analysis. We tested the numeric differences between species using the Mann–Whitney *U*-test. Additionally, we tested the hypothesis that the ladder web may be an adaptation to arboricolous life history and thus, that the LI and tree circumference would be negatively correlated within each size class. We analysed the data using SPSS, version 13.0.1 (SPSS Inc.). Where necessary, the probability levels were corrected by Bonferroni adjustment.

RESULTS

The technique of assigning spiders to size classes in the field is validated because the size classes that we inferred in the field correlated significantly with the measured lengths of patella + tibia I (*H. etruscilla*: $r = 0.943$, $N = 24$, $P < 0.001$; *H. multipuncta*: $r = 0.904$, $N = 23$, $P < 0.001$; *N. malabarensis*: $r = 0.788$, $N = 40$, $P < 0.001$).

The web parameters of both *Herennia* species did not differ significantly (Table 1); thus, we considered *H. multipuncta* webs to be representative of the genus (Fig. 2, Table 2). Typical small juvenile instar webs in both *Herennia* and *Nephilengys* were round orbs, which were built against tree bark (Fig. 1B, C), their webs had about the same levels of ladderness and HD (Fig. 2). Web size increase followed the predicted allometric shift towards ladder architecture (Fig. 2), which is more pronounced in *Herennia* (LI values rise from 0.9 to 5.5; Fig. 2) than in *Nephilengys*, where LI

Table 1. Probabilities (*P*) for interspecific *Herennia* differences (*Herennia multipuncta* versus *Herennia etruscilla*) in web parameters for the sparse data available for *H. etruscilla* ($N = 20$) (Mann–Whitney *U*-test)

Parameter	Stage 2	Stage 3	Stage 4	Stage 5	Stage 6	Stage 7	Stage 8
Tree circumference	NA	0.165	0.175	0.434	0.034	0.340	0.554
Ladder index	NA	0.299	0.636	0.235	0.724	NA	0.248
Hub displacement	NA	0.309	0.156	0.317	0.364	NA	0.739
Hub to bark	NA	0.490	0.129	1.000	1.000	NA	1.000
From ground	NA	0.933	0.195	0.373	0.289	NA	0.439

All *P*-values are nonsignificant. NA, not available.

Table 2. Probabilities for intergeneric differences (*Herennia multipuncta* versus *Nephilengys malabarensis*) in web parameters for each size class (Mann–Whitney *U*-test, chi-square analysis)

Parameter	Stage 2	Stage 3	Stage 4	Stage 5	Stage 6	Stage 7	Stage 8
Tree circumference	< 0.001	< 0.001	0.023	0.024	NA	NA	0.157
Ladder index	< 0.001	0.047	0.046	0.803	NA	NA	0.040
Side frames	0.646	< 0.001	0.436	0.232	NA	NA	0.047
Web shape	0.809	0.016	0.265	0.016	NA	NA	0.151
Presence/absence of hub cup	0.001	< 0.001	< 0.001	< 0.001	NA	NA	< 0.001
Presence/absence of pseudoradii	NA	NA	0.057	0.016	NA	NA	< 0.001
Presence/absence of retreat	< 0.001	< 0.001	< 0.001	< 0.001	NA	NA	< 0.001
Hub displacement	0.044	0.670	0.047	0.893	NA	NA	0.024
Hub to bark	< 0.001	< 0.001	< 0.001	0.007	NA	NA	0.044
From ground	< 0.001	0.005	0.014	1.000	NA	NA	0.558

Significant values are shown in bold. NA, not available.

reached a value slightly above 2 in the fifth instar (Fig. 2; interspecific differences are shown in Table 2).

In *Herennia*, web side frames changed from round in smaller size classes through subparallel to parallel in larger spiders ($\chi^2 = 27.3$, $N = 93$, $P = 0.007$) and the according web shape changed from planar to curved ($\chi^2 = 32.5$, $N = 95$, $P < 0.001$), with this curvature being convex and following the tree circumference (Fig. 1A). As reported previously, we confirmed the vertically parallel threads termed the pseudoradii (Fig. 1A) as apomorphic element of adult *Herennia* webs (Kuntner, 2005). We established their absence in webs of smaller size classes and their significantly increasing prevalence in larger size classes (Fig. 3). In *Nephilengys*, web side frames changed from round to subparallel ($\chi^2 = 37.2$, $N = 142$, $P < 0.001$), and web shape from planar to curved ($\chi^2 = 53.4$, $N = 144$, $P < 0.001$), although this curvature was concave (Fig. 1D). The webs of *Nephilengys* never possessed pseudoradii, nor a hub cup, but always a retreat connected to the hub (Fig. 1D), as reported previously for all *Nephilengys* species (Kuntner, 2007).

HD values did not show a significant ontogenetic increase in *Herennia* (Fig. 2), and not even in heavy-

bodied *Nephilengys*, with the exception of a pronounced increase between instars five and eight ($U = 1$, $N = 11$, $P = 0.009$; Fig. 2). There was an increase in distance from ground in the growing *Herennia* and *Nephilengys* webs (Fig. 2).

Both genera may utilize a similar habitat: forest patches under any canopy closure. The webs of all size classes in both genera always utilized the substrate (e.g. tree trunks) and thus were not aerial. Juvenile *Herennia* inhabited thinner trees compared to *Nephilengys* (Table 2). There were, however, also marked differences in the webs of *Herennia* and *Nephilengys*. The distance between the hub and the substrate decreased with spider size in *Herennia* (Fig. 2) and fell to zero, with the hub cup invariably touching bark in larger spiders (Figs 2, 3A). By contrast, in *Nephilengys*, the hub was positioned increasingly further away from the substrate in larger size classes (Fig. 2).

In *Herennia*, the correlation between LI and tree circumference was always negative, although this was only significant for the third- and fifth-instar data (second instar: $P = 0.827$; third: $P = 0.021$; fourth: $P = 0.413$; fifth: $P = 0.05$; seventh: $P = 0.239$; eighth: $P = 0.756$). In *Nephilengys*, the correlation was




Figure 3. Number of webs with or without hub cup and pseudoradiali in *Herennia multipunctata*: in contrast to juvenile webs, webs of larger instars exhibited the hub cup and pseudoradiali. *P*-values are probabilities for the difference between size classes (contingency tables, chi-square analysis).

never significant (second instar: $P = 0.756$; third: $P = 0.853$; fourth: $P = 0.730$).

DISCUSSION

The present study investigated the web architectures in most ontogenetic stages in two south-east Asian nephilid spiders belonging to the genera *Herennia* and *Nephilengys*. Both utilize tree trunks (Fig. 1), or sometimes other substrates, as websites. However, their web architectures differ considerably. Although small juvenile webs in both genera are round orbs (Fig. 1C), adult architectures become increasingly elongate, with extreme ladderness in *Herennia* (Fig. 1A). Additionally, larger instars in *Herennia* con-

struct their ladder webs closely adhering to the tree shape; thus, the web is convex, contains the unique pseudoradiali and a hub cup (Kuntner, 2005), and comes into contact with tree bark (Fig. 1A). On the other hand, larger instars in *Nephilengys* build wider, planar or concave webs on wider trees, with the lower part of the web extending away from the substrate. Their webs are more vertically asymmetric, with the hub displaced towards the top frame, and only come in touch with the substrate at the retreat (Fig. 1D) (Kuntner, 2007). These two different adult architectures make it possible for *Herennia* and *Nephilengys* to exploit different niches. Although all known *Herennia* are consistently arboricolous (Kuntner, 2005), *Nephilengys* species are found both on trees and around houses (Kuntner, 2007). The original micro-habitat for adult *Nephilengys* is the canopy, where large branches provide shelter for the spider's retreat, and ample aerial space for the capture web below. Such webs were well pre-adapted for the roofs of houses with a high substrate anchoring point suitable for the retreat, shelter from rain, and an abundance of flying insect prey around human dwellings.

As predicted for both *Herennia* and *Nephilengys*, the LI changed significantly with spider size/age, with larger spiders building more ladder-shaped webs (Fig. 2). This repeats the pattern observed in *Clitaetra* (Kuntner *et al.*, 2008b; Kuntner & Agnarsson, 2009), the sister clade to all other nephilids (Fig. 4) (Kuntner *et al.*, 2008a). The combined results point to an evolutionary strategy in arboricolous nephilid spiders (grade-lineage *Clitaetra–Herennia–Nephilengys*; Fig. 4), which enables the growing spider to enlarge its orb web allometrically such that it may remain on a given tree through ontogeny.

Several studies have compared web shapes using the same or equivalent quantifications of vertical elongation, or ladderness (Harmer & Framenau, 2008; Kuntner *et al.*, 2008b; Harmer, 2009; Harmer & Herberstein, 2009; Kuntner & Agnarsson, 2009). Surprisingly, however, the literature still lacks a clear definition of what constitutes a ladder web. Clearly, the extremely elongated aerial web spun by the neotropical araneid spider *Scoloderus* falls into this category (Eberhard, 1975) as do webs of phylogenetically scattered araneoid exemplars such as *Tylorida* (Robinson & Robinson, 1972) and *Herennia* (Robinson & Lubin, 1979; Kuntner, 2005). We treat as ladder webs those that reach and exceed twice the maximal length over maximal width (LI values above 2) and possess (sub)parallel, rather than round, side frames. Although arbitrary, this definition is based on the use in recent literature: in *Herennia*, the mean LI is between 1.5 (second instar) and 3.5 (eighth instar); in *Clitaetra*, the mean LI ranges from 1.9 in second instar to 4 in eighth instar; in adult *Telaprocera* the




Figure 4. Phylogenetic tree summarizing nephilid spider-web evolution, topology and events combined from Blackledge *et al.* (2009), Kuntner & Agnarsson (2009) Kuntner *et al.* (2008a, b) and the present study. The three species investigated in the present study are highlighted. Nephilid orb web shape is defined as 'ladder' in taxa showing a significant ontogenetic ladder index increase above the value $LI = 2$, and (sub)parallel, rather than round, side frames. Web position is 'arboricolous' (i.e. mostly found on tree trunks or utilizing substrate at least for the retreat) versus 'aerial' (i.e. not utilizing substrate). 'Hub eccentricity' is according to the known hub displacement increase (for details, see text).

mean LI is between 2.4 and 3.1 (Harmer, 2009). Thus, the ontogeny of most 'ladder-weavers' starts at $LI = 1-2$, and then increases to values above 2 in larger spiders. On the basis of this definition, also some *Nephilengys* instars build ladder webs (LI values slightly exceeding 2), although these are not as extremely elongated as the ladder webs in *Herennia* (LI values between 2 and 4). We consider that the real pattern would be even stronger in large *Nephilengys*,

which build webs high above the ground (see a trend shown in Fig. 2), and were thus largely inaccessible for our study (sixth- and seventh-instar data are particularly missing).

Phylogenetic evidence suggests that araneoid ladder webs evolved repeatedly (Blackledge *et al.*, 2009). Such web architecture may be a specialization to web site or prey type. The former is the case in the Australian araneid *Telaprocera* (Harmer & Fra-

menau, 2008; Harmer, 2009), and the latter in the neotropical *Scoloderus* (Eberhard, 1975; Stowe, 1978). Stowe (1978) showed that *Scoloderus* predominantly preys on moths; hence, the extreme elongation above the hub in its aerial web probably acts as an interception device. By contrast, *Telaprocera* do not specialize on a particular prey, and its substrate-web elongation is plastic and depends on the available space (Harmer & Herberstein, 2009). With a species-level phylogeny available (Kuntner *et al.*, 2008a; Kuntner & Coddington, 2009), nephilid spiders now offer an opportunity for a clade-wide web comparison (Fig. 4). No nephilid spider is known to specialize on a particular prey (Kuntner *et al.*, 2008b); thus, the adaptation to the web site appears plausible (Kuntner *et al.*, 2008b). Demonstrating an ontogenetic increase in ladderness in *Clitaetra*, Kuntner *et al.* (2008b) explained the nephilid ladder web as an adaptation to an arboricolous life history. However, if the *Telaprocera* levels of plasticity (Harmer & Herberstein, 2009) were to apply to nephilids, one would expect that the LI and tree circumference would be negatively correlated within each size class. In the present study, this was not the case for *Nephilengys*, although the results obtained for *Herennia* are ambiguous. In that case, the correlation was always negative, although this was only significant for the third- and fifth-instar data. Thus, although we cannot conclude whether ladderness in *Herennia* is a consequence of an evolutionary constraint versus plasticity, the former explains *Nephilengys* webs better.

Unlike in *C. irenae*, the hub relative position in *Herennia* and *Nephilengys* does not gradually shift towards the top web frame. Surprisingly, larger *Herennia* did not show significant increases in HD, but, to some degree, *Nephilengys* did. Such extreme asymmetry in adult females may have evolved in response to female gigantism in derived nephilids (Kuntner & Coddington, 2009) because gravity precludes efficient attacks upwards in heavy-bodied spiders (Masters & Moffat, 1983; Kuntner *et al.*, 2008b). Herberstein & Heiling (1999) showed that weight itself could explain vertical orb asymmetry in araneid spiders. Thus, HD ought to be less phylogenetically and ontogenetically constrained compared to other web features because the spider size itself is evolutionarily and ontogenetically labile. The lack of a trend in hub asymmetry increase in adult *Herennia* females may be an artefact of too few females in our sample. However, if such a pattern is real, it questions previous hypotheses of the effect of gravity (ap Rhisiart & Vollrath, 1994; Herberstein & Heiling, 1999) and calls for a more complex explanation, perhaps related to web-building costs (Coslovsky & Zschokke, 2009), ecological factors, or phylogenetic constraints.

Overall, the webs of *H. etruscilla* and *H. multipuncta* did not differ significantly (Table 1), suggesting uniformity of *Herennia* webs (Kuntner, 2005). Similarly, the web ontogeny in *N. malabarensis* resembled that documented in *N. cruentata* (Japyassu & Ades, 1998), which supports *Nephilengys* web conservatism (Kuntner, 2007). These data support the invoked nephilid intrageneric web uniformity in general (Kuntner & Agnarsson, 2009). Although nephilid web evolution spans four different web morphotypes (Kuntner *et al.*, 2008a), it appears to be slow and conserved within genera. Such slow changes are certainly not the rule in araneoid spiders. Eberhard *et al.* (2008a) showed rapid changes in web architectures, suggesting a remarkable evolutionary plasticity in the family Theridiidae.

Figure 4 summarizes our current understanding of nephilid web evolution. Regardless of whether nephilids are sister to all other araneoids (Fig. 4) (Kuntner *et al.*, 2008a) or nest within more derived araneoid clades (e.g. Blackledge *et al.*, 2009), the ancestral nephilid web form is reconstructed as a small ladder exemplified by *Clitaetra* (Kuntner, 2006; Kuntner *et al.*, 2008b; Kuntner & Agnarsson, 2009). Such a modification of the round orb web thus represents a reversal back to substrate-dependent architecture. Although each genus makes a different version, all behaviourally known *Clitaetra*, *Herennia*, and *Nephilengys* make substrate dependent webs, mostly ladders on trees. On the other hand, webs of *Nephila*, phylogenetically sister to *Nephilengys*, comprise aerial round orbs that only anchor to the substrate. The vertical elongation and substrate dependence in *Nephilengys* may be viewed as intermediate between substrate-ladders (e.g. *Herennia*) and aerial orbs (*Nephila*). Thus, the ancestor to the clade *Nephila* + *Nephilengys* probably tended to secondarily 'lose' the substrate. The complete freeing from the substrate, however, occurred in the ancestor to all extant *Nephila*, and this event must have represented an evolutionary repeat to the one at the root of the orbicularian tree (Fig. 4) (Blackledge *et al.*, 2009). In other words, if the orbicularian ancestor invented the aerial orb web, the *Nephila* ancestor, at a much less inclusive phylogenetic level, reinvented it.

ACKNOWLEDGEMENTS

We thank D. Li, D. Court, C. Koh, J. Woon, L. Yap, E. Tan, J. Koh, and P. Ng for field and/or logistical help, and three anonymous reviewers for their suggestions on how to improve the manuscript. This work was funded by the grants to MK from the European Community Sixth Framework Programme (a Marie Curie International Reintegration Grant MIRG-CT-

2005 036536), the Slovenian Research Agency (grant Z1-7082-0618) and the National University of Singapore (Raffles Museum Fellowship).

REFERENCES

- Agnarsson I.** 2003. Spider webs as habitat patches – the distribution of kleptoparasites (*Argyrodes*, Theridiidae) among host webs (*Nephila*, Tetragnathidae). *Journal of Arachnology* **31**: 344–349.
- Agnarsson I, Coddington JA.** 2006. Notes on web and web plasticity and description of the male of *Achaearanea hieroglyphica* (Mello-Leitao) (Araneae, Theridiidae). *Journal of Arachnology* **34**: 638–641.
- Agnarsson I, Dhinojwala A, Sahni V, Blackledge TA.** 2009. Spider silk as a novel high performance biomimetic muscle driven by humidity. *Journal of Experimental Biology* **212**: 1989–1993.
- ap Rhisiart A, Vollrath F.** 1994. Design features of the orb web of the spider, *Araneus diadematus*. *Behavioral Ecology* **5**: 280–287.
- Benjamin SP, Zschokke S.** 2003. Webs of theridiid spiders: construction, structure and evolution. *Biological Journal of the Linnean Society* **78**: 293–305.
- Benjamin SP, Zschokke S.** 2004. Homology, behaviour and spider webs: web construction behaviour of *Linyphia hortensis* and *L. triangularis* (Araneae: Linyphiidae) and its evolutionary significance. *Journal of Evolutionary Biology* **17**: 120–130.
- Blackledge TA.** 1998. Signal conflict in spider webs driven by predators and prey. *Proceedings of the Royal Society of London Series B, Biological Sciences* **265**: 1991–1996.
- Blackledge TA, Gillespie RG.** 2002. Estimation of capture areas of spider orb webs in relation to asymmetry. *Journal of Arachnology* **30**: 70–77.
- Blackledge TA, Gillespie RG.** 2004. Convergent evolution of behavior in an adaptive radiation of Hawaiian web-building spiders. *Proceedings of the National Academy of Sciences of the United States of America* **101**: 16228–16233.
- Blackledge TA, Scharff N, Coddington JA, Szuts T, Wenzel JW, Hayashi CY, Agnarsson I.** 2009. Reconstructing web evolution and spider diversification in the molecular era. *Proceedings of the National Academy of Sciences of the United States of America* **106**: 5229–5234.
- Blackledge TA, Summers AP, Hayashi CY.** 2005. Gumbooted lines in black widow cobwebs and the mechanical properties of spider capture silk. *Zoology* **108**: 41–46.
- Blackledge TA, Swindeman JE, Hayashi CY.** 2005. Quasistatic and continuous dynamic characterization of the mechanical properties of silk from the cobweb of the black widow spider *Latrodectus hesperus*. *Journal of Experimental Biology* **208**: 1937–1949.
- Blackledge TA, Zevenbergen JM.** 2006. Mesh width influences prey retention in spider orb webs. *Ethology* **112**: 1194–1201.
- Bond JE, Opell BD.** 1998. Testing adaptive radiation and key innovation hypotheses in spiders. *Evolution* **52**: 403–414.
- Coddington J.** 1986a. The monophyletic origin of the orb web. In: Shear WA, ed. *Spiders: webs, behavior and evolution*. Stanford, CA: Stanford University Press, 319–363.
- Coddington JA.** 1986b. The monophyletic origin of the orb web. In: Shear WA, ed. *Spiders: webs, behavior, and evolution*. Stanford, CA: Stanford University Press, 319–363.
- Coddington JA.** 1986c. Orb webs in ‘non-orb weaving’ ogre-faced spiders (Araneae: Dinopidae): a question of genealogy. *Cladistics* **2**: 53–67.
- Coddington JA.** 1989. Spinneret silk spigot morphology: Evidence for the monophyly of orb-weaving spiders, Cyrtophorinae (Araneidae), and the group Theridiidae plus Nesticidae. *Journal of Arachnology* **17**: 71–96.
- Coddington JA, Levi HW.** 1991. Systematics and evolution of spiders (Araneae). *Annual Review of Ecology and Systematics* **22**: 565–592.
- Coddington J, Sobrevila C.** 1987. Web manipulation and two stereotyped attack behaviors in the ogre-faced spider *Deinopis spinosus* Marx (Araneae, Deinopidae). *Journal of Arachnology* **15**: 213–225.
- Coslovsky M, Zschokke S.** 2009. Asymmetry in orb-webs: an adaptation to web building costs? *Journal of Insect Behavior* **22**: 29–38.
- Craig CL.** 2003. *Spider webs and silk: tracing evolution from molecules to genes to phenotypes*. New York, NY: Oxford University Press.
- Denny M.** 1976. Physical properties of spider silks and their role in design of orb-webs. *Journal of Experimental Biology* **65**: 483–506.
- Eberhard WG.** 1975. Inverted ladder orb web of *Scoloderus* sp. and intermediate orb of *Eustala* sp. Araneae – Araneidae. *Journal of Natural History* **9**: 93–106.
- Eberhard WG.** 1977. Rectangular orb webs of *Synotaxus* (Araneae-Theridiidae). *Journal of Natural History* **11**: 501–507.
- Eberhard WG.** 1981. Construction behaviour and the distribution of tensions in orb webs. *Bulletin of the British Arachnological Society* **5**: 189–204.
- Eberhard WG.** 1982. Behavioural characters for the higher classification of orb-weaving spiders. *Evolution* **36**: 1067–1095.
- Eberhard WG.** 1985. The ‘sawtoothed’ orb web of *Eustala* sp. (Araneae, Araneidae) with a discussion of ontogenetic changes in spiders’ web-building behavior. *Psyche (Cambridge)* **92**: 105–117.
- Eberhard WG.** 1986. Web building behavior of anapid, symphytognathid and mysmenid spiders (Araneae). *Journal of Arachnology* **14**: 339–356.
- Eberhard WG.** 1987. Hub construction by *Leucauge mariana* (Araneae, Araneidae). *Bulletin of the British Arachnological Society* **7**: 128–132.
- Eberhard WG.** 1990. Early stages of orb construction by *Philoponella vicina*, *Leucauge mariana*, and *Nephila clavipes* (Araneae, Uloboridae and Tetragnathidae), and their phylogenetic implications. *Journal of Arachnology* **18**: 205–234.
- Eberhard WG.** 1992. Web construction by *Modisimus* sp. (Araneae, Pholcidae). *Journal of Arachnology* **20**: 25–34.

- Eberhard WG.** 2007. Stabilimenta of *Philoponella vicina* (Araneae: Uloboridae) and *Gasteracantha cancriformis* (Araneae: Araneidae): evidence against a prey attractant function. *Biotropica* **39**: 216–220.
- Eberhard WG, Agnarsson I, Levi HW.** 2008a. Web forms and the phylogeny of theridiid spiders (Araneae: Theridiidae): chaos from order. *Systematics and biodiversity* **6**: 415–475.
- Eberhard WG, Barrantes G, Madrigal-Brenes R.** 2008b. Vestiges of an orb-weaving ancestor? The ‘biogenetic law’ and ontogenetic changes in the webs and building behavior of the black widow spider *Latrodectus geometricus* (Araneae Theridiidae). *Ethology Ecology & Evolution* **20**: 211–244.
- Forster LM, Forster RR.** 1985. A derivative of the orb web and its evolutionary significance. *New Zealand Journal of Zoology* **12**: 455–465.
- Garb J, DiMauro T, Vo V, Hayashi C.** 2006. Silk genes support the single origin of orb-webs. *Science* **312**: 1762.
- Griswold CE, Coddington JA, Hormiga G, Scharff N.** 1998. Phylogeny of the orb-web building spiders (Araneae, Orbiculariae: Deinopoidea, Araneoidea). *Zoological Journal of the Linnean Society* **123**: 1–99.
- Harmer AMT.** 2009. Elongated orb-webs of Australian ladder-web spiders (Araneidae: *Telaproceria*) and the significance of orb-web elongation. *Journal of Ethology* **27**: 453–460.
- Harmer AMT, Framenau JW.** 2008. *Telaproceria* (Araneae: Araneidae), a new genus of Australian orb-web spiders with highly elongated webs. *Zootaxa* **59**–80.
- Harmer AMT, Herberstein ME.** 2009. Taking it to extremes: what drives extreme web elongation in Australian ladder web spiders (Araneidae: *Telaproceria maudae*)? *Animal Behaviour* **78**: 499–504.
- Harvey MS, Austin AD, Adams M.** 2007. The systematics and biology of the spider genus *Nephila* (Araneae: Nephilidae) in the Australasian region. *Invertebrate Systematics* **21**: 407–451.
- Hayashi CY.** 2001. Convergent evolution of sequence elements in spider and insect silk proteins. *American Zoologist* **41**: 1468–1468.
- Hayashi C, Blackledge TA, Lewis R.** 2004. Molecular and mechanical characterization of aciniform silk: uniformity of iterated sequence modules in a novel member of the spider silk fibroin gene family. *Molecular Biology and Evolution* **21**: 1950–1959.
- Hayashi CY, Lewis RV.** 2000. Molecular architecture and evolution of a modular spider silk protein gene. *Science* **287**: 1477–1479.
- Herberstein ME, Heiling AM.** 1998. Does mesh height influence prey length in orb-web spiders (Araneae)? *European Journal of Entomology* **95**: 367–371.
- Herberstein ME, Heiling AM.** 1999. Asymmetry in spider orb webs: a result of physical constraints? *Animal Behaviour* **58**: 1241–1246.
- Higgins L.** 1992. Developmental changes in barrier web structure under different levels of predation risk in *Nephila clavipes* (Araneae, Tetragnathidae). *Journal of Insect Behavior* **5**: 635–655.
- Higgins L.** 2006. Quantitative shifts in orb-web investment during development in *Nephila clavipes* (Araneae, Nephilidae). *Journal of Arachnology* **34**: 374–386.
- Higgins LE, Buskirk RE.** 1992. A trap-building predator exhibits different tactics for different aspects of foraging behavior. *Animal Behaviour* **44**: 485–499.
- Japyassu HF, Ades C.** 1998. From complete orb to semi-orb webs: developmental transitions in the web of *Nephilengys cruentata* (Araneae: Tetragnathidae). *Behaviour* **135**: 931–956.
- Japyassu HF, Caires RA.** 2008. Hunting tactics in a cobweb spider (Araneae-Theridiidae) and the evolution of behavioral plasticity. *Journal of Insect Behavior* **21**: 258–284.
- Kuntner M.** 2005. A revision of *Herennia* (Araneae: Nephilidae: Nephilinae), the Australasian ‘coin spiders’. *Invertebrate Systematics* **19**: 391–436.
- Kuntner M.** 2006. Phylogenetic systematics of the Gondwanan nephilid spider lineage Clitaetinae (Araneae, Nephilidae). *Zoologica Scripta* **35**: 19–62.
- Kuntner M.** 2007. A monograph of *Nephilengys*, the pantropical ‘hermit spiders’ (Araneae, Nephilidae, Nephilinae). *Systematic Entomology* **32**: 95–135.
- Kuntner M, Agnarsson I.** 2009. Phylogeny accurately predicts behaviour in Indian Ocean *Clitaetra* spiders (Araneae: Nephilidae). *Invertebrate Systematics* **23**: 193–204.
- Kuntner M, Coddington JA.** 2009. Discovery of the largest orbweaving spider species: the evolution of gigantism in *Nephila*. *PLoS ONE* **4**: e7516.
- Kuntner M, Coddington JA, Hormiga G.** 2008a. Phylogeny of extant nephilid orb-weaving spiders (Araneae, Nephilidae): testing morphological and ethological homologies. *Cladistics* **24**: 147–217.
- Kuntner M, Haddad CR, Aljančić G, Blejec A.** 2008b. Ecology and web allometry of *Clitaetra irenae*, an arboreal African orb-weaving spider (Araneae, Araneoidea, Nephilidae). *Journal of Arachnology* **36**: 583–594.
- Li D.** 2005. Spiders that decorate their webs at higher frequency intercept more prey and grow faster. *Proceedings of the Royal Society of London Series B, Biological Sciences* **272**: 1753–1757.
- Masters WM, Moffat AJM.** 1983. A functional explanation of top-bottom asymmetry in vertical orbwebs. *Animal Behaviour* **31**: 1043–1046.
- Oppell BD, Schwend HS.** 2009. Adhesive efficiency of spider prey capture threads. *Zoology* **112**: 16–26.
- Robinson MH, Lubin YD.** 1979. Specialists and generalists: the ecology and behavior of some web-building spiders from Papua New Guinea, 2. *Psechrus argentatus* and *Fecenia* sp. (Araneae: Psechridae). *Pacific Insects* **21**: 133–164.
- Robinson MH, Robinson B.** 1972. The structure, possible function and origin of the remarkable ladder-web built by a New-Guinea orbweb spider (Araneae: Araneidae). *Journal of Natural History* **6**: 687–694.
- Rypstra AL.** 1982. Building a better insect trap – an experimental investigation of prey capture in a variety of spider webs. *Oecologia* **52**: 31–36.
- Scharff N, Coddington JA.** 1997. A phylogenetic analysis of the orb-weaving spider family Araneidae (Arachnida,

- Araneae). *Zoological Journal of the Linnean Society* **120**: 355–434.
- Schuck Paim C.** 2000. Orb-webs as extended-phenotypes: web design and size assessment in contests between *Nephilengys cruentata* females (Araneae: Tetragnathidae). *Behaviour* **137**: 1331–1347.
- Shear WA, ed.** 1986. *Spiders, webs, behavior, and evolution*. Stanford, CA: Stanford University Press.
- Stowe MK.** 1978. Observations of two nocturnal orbweavers that build specialized webs: *Scoloderus cordatus* and *Wixia ectypa* (Araneae: Araneidae). *Journal of Arachnology* **6**: 141–146.
- Toft S.** 1987. Microhabitat identity of two species of sheet-web spiders, field experimental demonstration. *Oecologia* **72**: 216–220.
- Tso IM.** 1996. Stabilimentum of the garden spider *Argiope trifasciata*: a possible prey attractant. *Animal Behaviour* **52**: 183–191.
- Tso IM.** 1998a. Isolated spider web stabilimentum attracts insects. *Behaviour* **135**: 311–319.
- Tso IM.** 1998b. Stabilimentum-decorated webs spun by *Cyclosa conica* (Araneae, Araneidae) trapped more insects than undecorated webs. *Journal of Arachnology* **26**: 101–105.
- Tso IM, Jiang SY, Blackledge TA.** 2007. Does the giant wood spider *Nephila pilipes* respond to prey variation by altering web or silk properties? *Ethology* **113**: 324–333.
- Vollrath F, Selden P.** 2007. The role of behavior in the evolution of spiders, silks, and webs. *Annual Review of Ecology Evolution and Systematics* **38**: 819–846.
- Wise DH.** 1993. *Spiders in ecological webs*. Cambridge: Cambridge University Press.
- Zschokke S.** 1993. The influence of the auxiliary spiral on the capture spiral in *Araneus diadematus* Clerck (Araneidae). *Bulletin of the British Arachnological Society* **9**: 169–173.
- Zschokke S.** 1999. Nomenclature of the orb-web. *Journal of Arachnology* **27**: 542–546.
- Zschokke S.** 2003. Spider-web silk from the Early Cretaceous. *Nature (London)* **424**: 636–637.
- Zschokke S, Henaut Y, Benjamin SP, Garcia-Ballinas JA.** 2006. Prey-capture strategies in sympatric web-building spiders. *Canadian Journal of Zoology – Revue Canadienne De Zoologie* **84**: 964–973.

APPENDIX

Table A1. Ecological and web data for the three nephilid spider species studied in Southeast Asia.

Species	Bark	Tree circumference (cm)	Canopy height (cm)	Size class	Width (cm)	Height (cm)	Hub size (mm)	Hub depth (mm)	Hub form (cm)	Web shape	Hub cup	Pseudoradii	Retreat	Web orientation				
<i>Herennia etruscula</i>																		
Medium	135.0	Open	8	20.0	80.0	22.5	0.0	103.0	Subparallel	Planar	Yes	Yes	No	No	East			
Medium	92.0	Partially	3	6.0	10.5	4.5	0.0	195.0	Subparallel	Planar	Yes	No	No	No	South west			
Rough	279.0	Partially	8	30.0	58.5	13.0	0.0	223.0	Subparallel	Curved	Yes	Yes	No	No	East			
Rough	142.0	Partially	6	18.5	61.0	29.0	0.0	72.0	Subparallel	Curved	Yes	Yes	No	No	North east			
Rough	114.0	Partially	6	19.0	46.0	26.0	0.0	112.0	Subparallel	Curved	Yes	Yes	No	No	North east			
Smooth	77.0	Partially	6	12.5	66.5	32.0	0.0	183.0	Parallel	Curved	Yes	Yes	No	No	North east			
Rough	147.0	Partially	3	11.0	24.5	13.5	0.0	162.0	Round	Planar	Yes	No	No	No	East			
Rough	117.0	Open	4	14.0	28.0	10.5	0.0	198.0	Subparallel	Curved	Yes	Yes	No	No	North east			
Medium	157.0	Open	5	20.0	35.0	16.0	0.0	205.0	Subparallel	Curved	Yes	Yes	No	No	East			
Medium	157.0	Open	4	10.0	31.0	13.0	0.0	225.0	Parallel	Curved	Yes	Yes	No	No	North east			
Rough	165.0	Closed	4	8.5	17.5	8.3	0.0	232.0	Subparallel	Curved	Yes	Yes	No	No	South			
Rough	155.0	Partially	4	7.5	22.5	12.0	0.0	185.0	Subparallel	Planar	Yes	Yes	No	No	West			
Smooth	83.0	Partially	5	5	32.0	14.0	0.0	225.0										
Smooth	83.0	Partially	5	9.0	32.0	7.5	0.0	130.0										
Smooth	83.0	Partially	6	5	32.0	4.3	8.0	154.0										
Rough	302.0	Partially	5	3	9.0	14.0	7.5	12.0	Subparallel	Planar	No	No	No	No	North east			
Rough	302.0	Partially	2	4.7	9.0	4.3	7.5	102.0	Parallel	Planar	No	No	No	No	North east			
Rough	82.0	Partially	3	7.0	18.0	12.5	5.5	154.0	Subparallel	Planar	Yes	Yes	No	No	North west			
Medium	113.0	Partially	3	4.4	12.5	14.0	2.0	203.0	Subparallel	Curved	No	No	No	No	East			
Medium	97.0	Partially	3	5.5	14.0	5.1	2.0	209.0	Round	Curved	Yes	Yes	No	No	South			
Rough	127.0	Partially	4	8.4	12.9	5.1	0.0	209.0	Round	Planar	No	No	No	No	North			
Rough	170.0	Partially	4	9.0	27.0	9.5	0.0	250.0	Subparallel	Planar	Yes	Yes	No	No	North east			
Medium	140.0	Partially	3	6.7	8.6	3.1	0.0	242.0	Round	Planar	Yes	Yes	No	No	South			
Rough	184.0	Partially	8	37.0	100.0	40.0	0.0	245.0	Subparallel	Curved	Yes	Yes	No	No	North west			
Rough	184.0	Partially	2	4.8	8.0	3.7	6.0	174.0	Round	Planar	No	No	No	No	South west			
Rough	184.0	Partially	2	6.5	8.3	3.5	0.0	186.0	Round	Curved	Yes	Yes	No	No	West			
Rough	184.0	Partially	3	6.8	14.7	8.2	0.0	161.0	Subparallel	Planar	Yes	Yes	No	No	South west			
Rough	184.0	Partially	3	4.3	14.2	7.2	0.0	144.0	Parallel	Planar	Yes	Yes	No	No	North west			
Rough	184.0	Partially	3	3.9	6.4	1.8	0.0	129.0	Round	Planar	Yes	Yes	No	No	West			
Rough	184.0	Partially	3	5.5	8.5	2.9	0.0	160.0	* Subparallel	Planar	Yes	Yes	No	No	South west			
Rough	184.0	Partially	4	6.0	15.0	6.5	0.0	114.0	Round	Planar	Yes	Yes	No	No	South west			
Rough	184.0	Partially	4	8.1	9.2	4.8	0.0	160.0	Subparallel	Planar	Yes	Yes	No	No	South west			
Rough	184.0	Partially	2	5.1	6.7	3.5	4.0	123.0	Round	Planar	No	No	No	No	North			
Rough	184.0	Partially	5	11.5	21.0	9.6	0.0	73.0	Subparallel	Curved	Yes	Yes	No	No	North west			
Rough	184.0	Partially	5	12.3	26.6	10.3	0.0	202.0	Subparallel	Curved	Yes	Yes	No	No	North west			
Rough	184.0	Partially	3	6.0	10.1	3.8	0.0	168.0	Subparallel	Planar	Yes	Yes	No	No	South			
Rough	184.0	Partially	4	7.0	9.0	6.0	0.0	217.0	Round	Planar	Yes	Yes	No	No	North west			
<i>Herennia multipunctata</i>																		

Table A1. Continued

Species	Barb	Drecreumference (cm)	Canopy	Size class	Width (cm)	Height (cm)	Hub height (cm)	Hub bar height (mm)	Hub top to ground (mm)	Front ground (cm)	Side frames	Web shape	Hub cup	Pseudoradii	Reorientat	Web orientation	
	Rough	184.0	Partially	3	5.1	5.0	2.5	0.0	188.0	Round	Planar	Yes	No	No	No	South west	
	Rough	184.0	Partially	3	7.0	8.6	5.1	0.0	200.0	Round	Planar	Yes	No	No	No	South west	
	Rough	184.0	Partially	3	6.4	11.2	8.0	0.0	209.0	Subparallel	Planar	Yes	No	No	No	South	
	Rough	184.0	Partially	3	4.9	12.0	5.6	0.0	219.0	Subparallel	Planar	Yes	No	No	No	South	
	Rough	184.0	Partially	8													
	Medium	100.0	Partially	2	3.1	3.6	1.6	0.0	200.0	Round	Planar	Yes	No	No	No	North east	
	Smooth	110.0	Partially	2	5.7	7.1	3.7	15.0	108.0	Round	Planar	No	No	No	No	South east	
	Smooth	110.0	Partially	2	6.5	9.5	4.5	5.0	179.0	Round	Planar	No	No	No	No	South east	
	Smooth	110.0	Partially	2	5.5	9.5	5.0	6.0	245.0	Subparallel	Planar	No	No	No	No	East	
	Smooth	110.0	Partially	2	5.0	12.5	5.1	2.0	138.0	Parallel	Planar	No	No	No	No	North	
	Rough	350.0	Partially	2	6.0	15.0	8.0	5.0	75.0	Subparallel	Planar	No	No	No	No	South east	
	Rough	350.0	Partially	2	5.0	8.5	4.5	15.0	135.0	Subparallel	Planar	No	No	No	No	North west	
	Smooth	108.0	Partially	2	6.5	12.0	7.0	12.0	160.0	Subparallel	Planar	No	No	No	No	South	
	Medium	75.0	Closed	2	4.0	9.5	4.2	6.0	160.0	Subparallel	Planar	No	No	No	No	South west	
	Medium	100.0	Partially	3	5.5	12.5	6.5	0.0	148.0	Subparallel	Curved	Yes	No	No	No	North east	
	Medium	100.0	Partially	3	6.5	8.0	4.0	0.0	166.0	Subparallel	Curved	Yes	No	No	No	North east	
	Medium	103.0	Partially	3	7.0	11.0	6.0	0.0	240.0	Subparallel	Curved	Yes	No	No	No	East	
	Smooth	110.0	Partially	3	10.0	16.0	7.5	18.0	142.0	Round	Planar	No	No	No	No	South east	
	Rough	350.0	Partially	3	7.4	10.0	4.5	0.0	62.0	Round	Planar	Yes	No	No	No	North east	
	Rough	350.0	Partially	3	5.0	6.5	3.0	0.0	163.0	Round	Curved	Yes	No	No	No	North east	
	Rough	350.0	Partially	3	7.0	9.0	4.5	0.0	163.0	Round	Curved	Yes	No	No	No	North east	
	Medium	100.0	Partially	4	9.0	23.5	10.5	11.0	220.0	Subparallel	Planar	No	No	No	No	North east	
	Medium	100.0	Partially	4	7.0	18.0	9.0	9.0	214.0	Subparallel	Planar	No	No	No	No	East	
	Rough	109.0	Partially	4	5.5	7.0	4.0	0.0	137.0	Round	Curved	Yes	No	No	No	South west	
	Medium	340.0	Closed	4	8.0	17.0	10.0	0.0	192.0	Subparallel	Curved	Yes	No	No	No	North	
	Smooth	75.0	Open	5	11.0	39.0	16.5	0.0	202.0	Parallel	Curved	Yes	Yes	No	No	South	
	Smooth	75.0	Open	6	10.0	55.0	18.0	0.0	105.0	Parallel	Curved	Yes	No	No	No	South	
	Medium	210.0	Closed	2	8.0	7.0	4.0	5.0	78.0	Round	Planar	No	No	No	No	North west	
	Medium	210.0	Closed	3	6.8	14.0	5.7	6.0	100.0	Subparallel	Planar	No	No	No	No	West	
	Medium	210.0	Closed	2	6.2	11.5	4.0	3.0	137.0	Subparallel	Planar	No	No	No	No	South west	
	Medium	210.0	Closed	3	5.3	13.2	6.2	6.0	62.0	Round	Subparallel	Planar	No	No	No	No	South west
	Medium	210.0	Closed	2	8.1	11.1	5.7	3.0	156.0	Subparallel	Round	Curved	Yes	No	No	No	South east
	Medium	210.0	Closed	2	4.0	11.0	6.0	5.0	211.0	Parallel	Planar	No	No	No	No	South east	
	Medium	210.0	Closed	2	2.3	7.1	2.9	0.0	173.0	Parallel	Curved	Yes	No	No	No	East	
	Medium	210.0	Closed	2	6.2	11.5	5.7	6.0	199.0	Subparallel	Planar	No	No	No	No	South west	
	Medium	210.0	Closed	3	6.0	11.0	5.5	0.0	225.0	Subparallel	Planar	Yes	No	No	No	South west	
	Medium	210.0	Closed	3	6.0	8.5	5.8	0.0	147.0	Subparallel	Round	Curved	Yes	No	No	No	South east
	Medium	210.0	Closed	2	7.6	8.2	3.6	3.0	177.0	Subparallel	Planar	Yes	No	No	No	South east	
	Medium	210.0	Closed	3	5.0	11.0	6.0	6.0	172.0	Subparallel	Planar	Yes	No	No	No	East	
	Medium	210.0	Closed	2	4.7	9.2	5.0	0.0	206.0	Subparallel	Planar	Yes	No	No	No	South west	
	Medium	210.0	Closed	3	7.1	12.6	6.0	9.0	175.0	Subparallel	Planar	No	No	No	No	North west	
	Medium	120.0	Partially	3	6.0	11.0	5.5	0.0	180.0	Parallel	Curved	Yes	No	No	No	North west	
	Medium	120.0	Partially	3	7.1	12.5	6.5	2.0	145.0	Subparallel	Planar	No	No	No	No	South west	
	Medium	120.0	Partially	2	6.0	10.8	5.7	3.0	146.0	Subparallel	Planar	No	No	No	No	South west	
	Medium	120.0	Partially	2	4.6	10.1	4.4	4.0	190.0	Subparallel	Planar	No	No	No	No	South west	

Medium	127.0	Partially	4	8.0	23.4	9.0	218.0	3.0	No
Medium	162.0	Partially	2	9.5	13.0	5.5	Subparallel	No	No
Rough	150.0	Closed	3	10.5	15.0	6.5	Subparallel	No	No
Rough	150.0	Closed	3	6.5	14.0	6.2	Round	No	No
Medium	227.0	Closed	2	6.0	7.5	3.0	Round	No	No
Medium	227.0	Closed	2	7.5	10.5	6.0	Round	No	No
Medium	227.0	Closed	2	7.0	10.0	5.0	190.0	No	No
Medium	227.0	Closed	2	7.1	9.0	4.0	230.0	No	No
Medium	227.0	Closed	2	5.5	7.5	3.7	180.0	No	No
Medium	227.0	Closed	3	11.5	17.0	7.2	175.0	No	No
Medium	227.0	Closed	3	6.0	12.5	6.9	12.0	No	No
Medium	227.0	Closed	3	5.5	13.2	5.6	150.0	No	No
Medium	157.0	Partially	2	6.7	8.0	3.9	125.0	No	No
Medium	157.0	Partially	2	6.0	8.1	3.5	100.0	No	No
Smooth	94.0	Partially	2	5.3	9.5	4.1	9.0	No	No
Medium	135.0	Partially	2	5.6	8.5	5.0	11.0	No	No
Medium	135.0	Partially	2	4.5	8.5	3.7	120.0	No	No
Medium	120.0	Closed	2	5.0	10.0	4.0	137.0	No	No
Medium	296.0	Closed	2	8.0	10.4	4.4	132.0	No	No
Medium	95.0	Partially	3	9.0	12.5	5.5	183.0	No	No
Medium	157.0	Partially	3	6.1	12.5	6.5	12.0	No	No
Smooth	94.0	Partially	3	6.6	7.6	8.5	216.0	No	No
Medium	122.0	Partially	3	9.5	21.5	10.0	167.0	No	No
Medium	122.0	Partially	4	10.0	21.0	10.0	200.0	No	No
Smooth	73.0	Partially	3	6.5	16.0	7.3	177.0	No	No
Smooth	95.0	Partially	3	3.6	18.0	8.0	200.0	No	No
Smooth	44.0	Partially	6	7.5	26.0	16.0	126.0	No	No
Medium	102.0	Partially	7	16.0	72.0	21.0	115.0	No	No
Medium	100.0	Partially	3	8.0	15.0	5.5	240.0	No	No
Medium	60.0	Partially	3	5.8	15.0	4.0	132.0	No	No
Medium	67.0	Partially	3	4.9	19.1	9.8	180.0	No	No
Medium	93.0	Partially	4	8.0	31.0	15.0	105.0	No	No
Medium	143.0	Partially	5	11.0	30.0	14.0	260.0	No	No
Medium	70.0	Partially	6	20.0	50.0	15.0	245.0	No	No
Medium	124.0	Partially	4	8.5	19.0	9.5	230.0	No	No
Medium	112.0	Partially	7	5.8	26.0	8.0	200.0	No	No
Smooth	73.0	Partially	7	15.0	53.0	22.0	105.0	No	No
Rough	218.0	Closed	5	12.5	29.0	13.0	9.0	No	No
Medium	340.0	Closed	3	10.0	22.5	11.5	150.0	No	No
Medium	236.0	Closed	4	15.0	23.0	13.0	103.0	No	No
Medium	340.0	Closed	5	16.0	38.0	16.0	102.0	No	No
Rough	400.0	Closed	5	21.0	51.0	24.5	110.0	No	No
Rough	121.0	Open	8	25.0	32.0	0.0	205.0	No	No
Smooth	67.0	Partially	3	10.3	29.2	3.5	130.0	No	No
Rough	197.0	Partially	4	12.1	22.0	10.0	138.0	No	No
Rough	197.0	Closed	2	10.0	9.5	3.5	124.0	No	No
Rough	197.0	Closed	3	16.0	24.5	11.0	141.0	No	No
Nephilengys malabarensis		Closed	3	15.5	28.7	13.0	187.0	No	No
Rough	197.0	Closed	3	16.0	25.5	9.0	190.0	No	No
Rough	197.0	Closed	3	8.0	14.5	6.0	105.0	No	No
Medium	303.0	Partially	2	6.2	9.1	4.4	212.0	No	No
Medium	303.0	Partially	2	7.8	7.2	3.2	180.0	No	No
							163.0	No	No

Table A1. Continued

Species	Barak	Tree circumference (cm)	Canopy size class	Width (cm)	Height (cm)	Hub to top (mm)	Hub to bark (mm)	Frontal form	Side frames	Hub cup	Web shape	Pseudodradii	Retreat	Web orientation
Medium	303.0	Partially	2	10.5	3.7	85.0	106.0	Round	Planar	No	Central	No	South west	
Medium	303.0	Partially	2	8.1	9.5	2.5	48.0	Round	Planar	No	Central	No	South	
Medium	303.0	Partially	2	6.0	7.1	1.5	28.0	Subparallel	Planar	No	Central	No	West	
Medium	303.0	Partially	2	7.5	11.8	3.6	28.0	Round	Planar	No	Central	No	Central	
Medium	303.0	Partially	2	7.5	14.0	4.2	16.0	60.0	Planar	No	Central	No	North west	
Medium	303.0	Partially	2	8.5	11.0	6.0	12.0	150.0	Planar	No	Central	No	North	
Medium	303.0	Partially	2	7.7	10.5	4.7	15.0	117.0	Planar	No	Central	No	North	
Medium	303.0	Partially	2	6.7	8.5	4.1	46.0	64.0	Round	Planar	No	Central	North east	
Medium	303.0	Partially	2	5.0	5.5	2.5	41.0	82.0	Planar	No	Central	No	North east	
Medium	303.0	Partially	2	5.0	9.0	5.0	36.0	106.0	Subparallel	Planar	No	Central	No	North east
Medium	303.0	Partially	2	8.0	11.5	6.0	14.0	144.0	Planar	No	Central	No	North east	
Medium	292.0	Partially	2	6.5	8.0	3.7	17.0	73.0	Planar	No	Central	No	North west	
Medium	292.0	Partially	2	7.5	12.5	5.0	36.0	77.0	Subparallel	Planar	No	Central	No	South west
Medium	292.0	Partially	2	9.5	12.0	5.0	17.0	172.0	Round	Planar	No	Central	No	South west
Medium	292.0	Partially	2	6.5	8.0	3.2	21.0	120.0	Round	Planar	No	Central	No	South west
Medium	292.0	Partially	2	7.7	8.6	4.2	35.0	110.0	Planar	No	Central	No	South west	
Medium	292.0	Partially	2	7.8	10.5	3.6	37.0	78.0	Planar	No	Central	No	South east	
Medium	66.0	Partially	2	5.2	6.8	2.3	31.0	190.0	Round	Planar	No	Central	No	North
Medium	66.0	Partially	2	7.0	9.0	4.5	25.0	216.0	Round	Planar	No	Central	No	North west
Medium	66.0	Partially	2	5.5	9.5	4.0	28.0	105.0	Subparallel	Planar	No	Central	No	South east
Smooth	77.0	Partially	2	7.5	10.0	5.0	16.0	75.0	Round	Planar	No	Central	No	South east
Smooth	77.0	Partially	2	7.1	7.2	3.9	20.0	125.0	Round	Planar	No	Central	No	South east
Medium	197.0	Partially	2	8.1	9.2	3.0	24.0	50.0	Round	Planar	No	Central	No	North
Medium	197.0	Partially	2	3.5	8.0	3.0	15.0	45.0	Subparallel	Planar	No	Central	No	North west
Medium	197.0	Partially	2	6.2	7.5	4.5	20.0	36.0	Round	Planar	No	Central	No	South east
Medium	273.0	Partially	2	6.5	9.0	3.5	50.0	117.0	Round	Planar	No	Central	No	North
Rough	183.0	Partially	2	7.0	10.0	5.2	23.0	73.0	Round	Planar	No	Central	No	South west
Rough	183.0	Partially	2	5.5	5.0	1.5	25.0	22.0	Round	Planar	No	Central	No	East
Rough	120.0	Partially	2	7.0	9.5	3.0	16.0	38.0	Round	Planar	No	Central	No	South east
Rough	315.0	Partially	2	5.5	8.5	3.3	10.0	105.0	Round	Planar	No	Central	No	North
Rough	315.0	Partially	2	7.5	9.5	4.0	65.0	95.0	Round	Planar	No	Central	No	North west
Rough	315.0	Partially	2	6.0	7.2	4.1	40.0	90.0	Round	Planar	No	Central	No	South west
Rough	120.0	Partially	2	7.0	8.0	2.5	25.0	167.0	Subparallel	Planar	No	Central	No	East
Rough	120.0	Partially	2	7.2	15.0	6.0	8.0	110.0	Round	Planar	No	Central	No	North
Rough	315.0	Partially	2	7.0	9.5	5.0	80.0	43.0	Round	Planar	No	Central	No	South west
Rough	120.0	Partially	2	6.0	8.0	4.6	40.0	20.0	Round	Planar	No	Central	No	South
Rough	325.0	Partially	2	6.5	7.5	3.5	30.0	105.0	Round	Planar	No	Central	No	North
Rough	325.0	Partially	2	9.0	10.7	5.7	15.0	74.0	Round	Planar	No	Central	No	East
Rough	325.0	Partially	2	7.0	13.5	6.0	13.0	68.0	Round	Planar	No	Central	No	North west
Rough	202.0	Partially	2	6.5	6.5	3.1	40.0	64.0	Round	Planar	No	Central	No	South
Rough	202.0	Partially	2	10.3	13.0	5.2	45.0	48.0	Round	Planar	No	Central	No	Central

Rough	202.0	Partially	2	5.5	5.0	2.5	20.0	44.0		No	Central	South east	
Rough	202.0	Partially	2	7.0	7.4	2.4	31.0	96.0		Planar	Planar	South east	
Medium	150.0	Partially	2	9.0	7.5	3.5	28.0	44.0		Planar	Planar	East	
Medium	150.0	Partially	2	8.1	10.3	4.9	24.0	122.0		Planar	Planar	East	
Medium	303.0	Partially	3	11.1	14.0	6.0	34.0	137.0		Planar	Planar	South west	
Medium	303.0	Partially	3	11.6	6.5	7.6	26.0	187.0		Planar	Planar	South	
Medium	303.0	Partially	3	10.0	11.5	5.5	17.0	183.0		Planar	Planar	North	
Medium	303.0	Partially	3	7.5	8.0	3.4	25.0	126.0		Planar	Planar	North east	
Medium	303.0	Partially	3	8.2	12.5	6.5	13.0	194.0		Planar	Planar	East	
Medium	303.0	Partially	3	10.3	11.5	5.6	20.0	163.0		Planar	Planar	South east	
Medium	292.0	Partially	3	11.0	12.5	6.9	16.0	88.0		Planar	Planar	North west	
Medium	292.0	Partially	3	10.0	13.0	5.5	44.0	93.0		Planar	Planar	South west	
Medium	292.0	Partially	3	9.2	15.0	7.0	25.0	167.0		Planar	Planar	South east	
Medium	66.0	Partially	3	8.0	13.0	8.3	29.0	191.0		Planar	Planar	West	
Medium	66.0	Partially	3	10.5	15.0	9.0	25.0	225.0		Planar	Planar	Lateral	
Medium	273.0	Partially	3	10.0	17.5	9.0	17.0	95.0		Planar	Planar	North	
Medium	273.0	Partially	3	11.8	18.0	9.9	130.0	16.0		Planar	Planar	East	
Medium	273.0	Partially	3	9.5	20.5	11.5	25.0	45.0		Planar	Planar	South east	
Rough	315.0	Partially	3	11.0	26.0	16.5	20.0	132.0		Subparallel	Subparallel	South east	
Rough	315.0	Partially	3	5.7	12.0	4.0	14.0	106.0		Parallel	Parallel	West	
Rough	120.0	Partially	3	9.5	16.0	8.0	16.0	135.0		Round	Round	North west	
Rough	325.0	Partially	3	10.0	13.5	5.0	12.0	35.0		Round	Round	South east	
Rough	202.0	Partially	3	6.5	14.5	6.3	22.0	130.0		Round	Round	West	
Rough	202.0	Partially	3	10.0	10.5	4.0	25.0	109.0		Subparallel	Subparallel	West	
Medium	150.0	Partially	3	13.2	17.0	12.0	42.0	75.0		Round	Round	South	
Medium	303.0	Partially	3	9.0	13.0	4.5	30.0	83.0		Subparallel	Subparallel	East	
Medium	303.0	Partially	4	13.0	26.0	13.0	30.0	177.0		Round	Round	North west	
Medium	197.0	Partially	4	13.0	25.0	8.0	31.0	143.0		Subparallel	Subparallel	South	
Medium	185.0	Partially	4	10.0	22.5	8.0	22.0	135.0		Subparallel	Subparallel	North	
Medium	350.0	Partially	4	12.0	20.5	9.0	45.0	25.0		Subparallel	Subparallel	North	
Rough	202.0	Partially	4	16.0	27.0	10.5	30.0	110.0		Subparallel	Subparallel	South	
Medium	150.0	Partially	4	12.0	10.5	3.5	26.0	70.0		Subparallel	Subparallel	East	
Rough	202.0	Partially	4	11.5	23.0	11.0	42.0	56.0		Subparallel	Subparallel	South	
Rough	202.0	Partially	5	11.0	10.0	6.5	55.0	68.0		Subparallel	Subparallel	West	
Medium	350.0	Partially	5	27.0	31.0	14.0	52.0	76.0		Subparallel	Subparallel	North west	
Rough	150.0	Partially	5	22.0	11.0	4.5	50.0	170.0		Round	Round	North west	
Medium	350.0	Partially	5	22.0	15.0	4.5	42.0	180.0		Planar	Planar	South west	
Rough	202.0	Partially	5	7.1	9.0	4.5	42.0	122.0		Subparallel	Subparallel	South east	
Rough	202.0	Partially	8	16.0	33.0	2.0	40.0	170.0		Subparallel	Subparallel	South east	
Medium	350.0	Partially	8	37.0	87.0	1.0	35.0	222.0		Curved	Curved	South east	
Medium	350.0	Partially	4	17.5	37.0	4.0	60.0	222.0		Subparallel	Subparallel	West	
Rough	218.0	Partially	5	52.0	52.0	4.5	60.0	225.0		Planar	Planar	Central	
Rough	218.0	Partially	8	50.0	82.0	3.0	55.0	105.0		Planar	Planar	South	
Rough	218.0	Partially	8	38.0	71.0	3.0	90.0	122.0		Curved	Curved	North west	
Rough	218.0	Closed	2	6.5	10.5	6.0	30.0	185.0		Curved	Curved	South east	
Rough	218.0	Closed	2	6.2	14.2	6.0	70.0	194.0		Subparallel	Subparallel	South east	
Rough	218.0	Closed	2	5.9	8.0	3.9	60.0	178.0		Subparallel	Subparallel	West	
Rough	218.0	Closed	2	5.0	6.6	3.5	105.0	225.0		Planar	Planar	South	
Rough	218.0	Closed	2	4.0	12.0	5.0	200.0	108.0		Parallel	Parallel	North west	
Rough	218.0	Closed	2	6.5	10.5	6.0	108.0	108.0		Subparallel	Subparallel	South east	
Rough	218.0	Closed	2	6.2	14.2	6.0	108.0	193.0		Planar	Planar	North east	

Table A1. Continued

Species	Barak extremity (cm)	Canopy extremity (cm)	Size class	Width (cm)	Height (cm)	Top to hub (cm)	Hub to barb (mm)	Front ground (cm)	Side frames	Web shape	Hub cup	Pseudoradii	Retreat	Web orientation	
Rough	218.0	Closed	2	7.0	7.0	3.2	200.0	Round	Planar	No	No	Central	Central	North west	North west
Rough	218.0	Closed	2	7.6	10.2	4.5	214.0	Round	Planar	No	No	Central	Central	North west	South
Medium	250.0	Closed	2	6.0	7.5	2.2	190.0	Round	Planar	No	No	Central	Central	South	South
Medium	250.0	Closed	2	6.0	8.4	3.1	180.0	Round	Planar	No	No	Central	Central	South	South
Medium	250.0	Closed	2	5.3	9.1	3.0	180.0	Subparallel	Planar	No	No	Central	Central	South east	South east
Medium	250.0	Closed	2	5.0	6.0	2.1	218.0	Subparallel	Planar	No	No	Central	Central	South east	South east
Medium	250.0	Closed	2	6.0	9.0	3.8	218.0	Subparallel	Planar	No	No	Central	Central	South east	South east
Medium	250.0	Closed	2	4.2	5.6	1.8	218.0	Round	Planar	No	No	Central	Central	South east	East
Medium	250.0	Closed	2	6.9	6.0	2.0	173.0	Round	Planar	No	No	Central	Central	East	East
Medium	250.0	Closed	2	5.5	12.0	4.0	108.0	Subparallel	Planar	No	No	Central	Central	East	East
Medium	250.0	Closed	2	7.0	8.5	3.7	160.0	Round	Planar	No	No	Central	Central	North west	North west
Medium	250.0	Closed	2	6.0	9.0	4.8	68.0	Round	Planar	No	No	Central	Central	North west	North west
Medium	250.0	Closed	2	5.0	6.3	2.9	180.0	Round	Planar	No	No	Central	Central	North west	North west
Medium	102.0	Closed	2	11.0	12.5	6.0	96.0	Round	Planar	No	No	Central	Central	North	North
Medium	102.0	Closed	2	8.5	12.5	4.5	50.0	Round	Planar	No	No	Central	Central	North	North
Medium	102.0	Closed	2	9.8	5.9	7.5	95.0	Subparallel	Planar	No	No	Central	Central	South east	South east
Medium	102.0	Closed	2	7.5	10.5	4.7	90.0	Round	Planar	No	No	Central	Central	West	West
Medium	184.0	Closed	2	7.4	8.5	5.3	60.0	Round	Planar	No	No	Central	Central	North west	North west
Medium	184.0	Closed	2	5.6	7.4	2.8	126.0	Round	Planar	No	No	Central	Central	South east	South east
Medium	265.0	Partially	2	12.0	18.0	8.0	76.0	Round	Planar	No	No	Central	Central	North	North
Medium	265.0	Partially	2	8.5	8.0	3.5	106.0	Round	Planar	No	No	Central	Central	West	West
Medium	265.0	Partially	2	10.0	12.4	6.0	78.0	Round	Planar	No	No	Central	Central	South	South
Medium	265.0	Partially	2	9.1	12.5	5.6	88.0	Round	Planar	No	No	Central	Central	South	South
Medium	265.0	Partially	2	6.0	6.0	2.2	71.0	Round	Planar	No	No	Central	Central	West	West
Medium	265.0	Partially	2	8.0	10.5	3.0	100.0	Subparallel	Planar	No	No	Central	Central	East	East
Rough	218.0	Closed	3	8.0	8.5	3.0	110.0	Round	Planar	No	No	Central	Central	South	South
Rough	218.0	Closed	3	6.0	15.0	7.5	183.0	Subparallel	Planar	No	No	Central	Central	North west	North west
Medium	250.0	Partially	2	6.5	10.2	5.6	107.0	Round	Planar	No	No	Central	Central	North west	North west
Medium	184.0	Closed	3	10.0	15.0	7.0	123.0	Subparallel	Planar	No	No	Central	Central	East	East
Medium	265.0	Partially	3	11.3	17.0	8.0	55.0	Round	Planar	No	No	Central	Central	East	East
Medium	265.0	Partially	3	12.0	21.0	13.0	228.0	Round	Planar	No	No	Central	Central	East	East
Medium	265.0	Partially	3	14.0	18.0	8.3	210.0	Round	Planar	No	No	Central	Central	North west	North west
Rough	102.0	Closed	4	11.6	18.0	12.0	190.0	Subparallel	Planar	No	No	Central	Central	North	North
Medium	265.0	Partially	4	9.6	12.0	4.0	78.0	Round	Planar	No	No	Central	Central	West	West
Medium	250.0	Closed	8	44.0	67.0	0.0	164.0	Subparallel	Planar	No	No	Central	Central	West	West